

On the Hanna Neumann Conjecture

Toshiaki Jitsukawa Bilal Khan Alexei G. Myasnikov

Abstract

The **Hanna Neumann conjecture** states that if F is a free group, then for all nontrivial finitely generated subgroups $H, K \leq F$,

$$\text{rank}(H \cap K) - 1 \leq [\text{rank}(H) - 1][\text{rank}(K) - 1]$$

Where most papers to date have considered a direct graph theoretic interpretation of the conjecture, here we consider the use of monomorphisms. We illustrate the effectiveness of this approach with two results. First, we show that for any finitely generated groups $H, K \leq F$ either the pair H, K or the pair H^-, K satisfy the Hanna Neumann conjecture; here $-$ denotes the automorphism which sends each generator of F to its inverse. Next, using particular monomorphisms from F to F_2 , we obtain that if the Hanna Neumann conjecture is false then there is a counterexample $H, K \leq F_2$ having the additional property that all the branch vertices in the foldings of H and K are of degree 3, and all degree 3 vertices have the same local structure or “type”.

1 Introduction

H. Neumann proved in [12] that any nontrivial subgroups $H, K \leq_{f.g.} F$ (finitely generated) must satisfy

$$\text{rank}(H \cap K) - 1 \leq 2[\text{rank}(H) - 1][\text{rank}(K) - 1], \quad (1)$$

and so improved Howson’s earlier result [5] that $H \cap K$ is finitely generated. The stronger assertion obtained by omitting the factor of 2 in (1) has come to be known as the Hanna Neumann conjecture. In [1], R. Burns improved H. Neumann’s bound by showing that

$$\begin{aligned} \text{rank}(H \cap K) - 1 &\leq 2[\text{rank}(H) - 1][\text{rank}(K) - 1] \\ &\quad - \min(\text{rank}(H) - 1, \text{rank}(K) - 1). \end{aligned}$$

In 1983, J. Stallings introduced the notion of a *folding* and showed how to apply these objects to the study of subgroups of free groups [16]. Stallings’s approach was applied by S. Gersten in [4] to solve certain special cases of the conjecture, and similar techniques were developed over a sequence of papers by W. Imrich [7, 6], P. Nickolas [14], and B. Servatius [15] who gave alternate

proofs of Burns' bound and resolved special cases of the conjecture. In 1989, W. Neumann showed that the conjecture is true "with probability 1" for randomly chosen subgroups of free groups [13], and proposed a stronger form of the conjecture. In 1992, G. Tardos proved in [17] that the conjecture is true if one of the two subgroups has rank 2. Then, in 1994, W. Dicks showed that the strong Hanna Neumann conjecture is equivalent to a conjecture on bipartite graphs, which he termed the Amalgamated Graph conjecture [2]. In 1996, G. Tardos used Dicks' method to give the first new bound for the general case in [18], where he proved that for any $H, K \leq F$ with $\text{rank}(H), \text{rank}(K) \geq 3$,

$$\begin{aligned} \text{rank}(H \cap K) - 1 &\leq 2[\text{rank}(H) - 1][\text{rank}(K) - 1] \\ &\quad - [\text{rank}(H) - 1] - [\text{rank}(K) - 1] \end{aligned}$$

Since then, W. Dicks and E. Formanek [3] proved that

$$\begin{aligned} \text{rank}(H \cap K) - 1 &\leq [\text{rank}(H) - 1][\text{rank}(K) - 1] + \\ &\quad \max\{\text{rank}(H) - 3, 0\} \max\{\text{rank}(K) - 3, 0\}, \end{aligned}$$

This resolved the conjecture for the case when one of the subgroups has rank at most 3.

The conjecture was also recently solved in the special case when one of the two groups, say H , has a generating set consisting of positive words (i.e. a set of words in which no generator of F has negative exponent). Specifically, it was shown by J. Meakin and P. Weil [11], and independently by B. Khan [9] that if there is some automorphism of F which carries a generating set of H to a set of positive words, then the conjecture holds for H and any nontrivial $K \leq_{f.g.} F$.

Recall that an automorphism σ of $F(X)$ is called *length-preserving* if $\forall u \in F$, $|u^\sigma| = |u|$, i.e. $(X^\pm)^\sigma = X^\pm$ where $X^\pm = X \cup X^{-1}$. In section 3, we shall prove the following two theorems:

Theorem 1. *Let $F_2 = F(a, b)$. Take $\sigma \in \text{Aut}(F_2)$ to be any length-preserving automorphism having no non-trivial fixed points, and let τ be any monomorphism*

$$\tau : \begin{array}{ll} a & \mapsto aw_aa \\ b & \mapsto bw_b b, \end{array}$$

where $w_a, w_b \in F_2$ are arbitrary elements for which the words aw_aa and $bw_b b$ are reduced as written. Then for all nontrivial $H, K \leq_{f.g.} F_2$, either the pair H, K or the pair $H^{\tau\sigma}, K$ satisfy the conjecture.

Theorem 2. *Let $F = F(X)$ and $-$ be the automorphism given by $x \mapsto x^{-1}$ (for each $x \in X$). Then for all nontrivial $H, K \leq_{f.g.} F$, either the pair H, K or the pair H^-, K satisfy the conjecture.*

Recall that given $H = \langle w_1, \dots, w_n \rangle \leq F$, one may determine the associated Stallings' folding $\Gamma_H = (V_H, E_H)$, by the following constructive procedure (see

[16]): Construct n directed cycles $c_1 = (V_1, E_1), \dots, c_n = (V_n, E_n)$, where $|V_i| = |w_i|$. Then pick one vertex from each of the cycles, and identify this subset of vertices, denoting the resulting vertex 1_H . Label the edges of cycle c_i by successive letters of w_i , starting at vertex 1_H . Finally, repeatedly identify pairs of edges e, e' for which

$$\text{label}(e) = \text{label}(e') \wedge [\text{head}(e) = \text{head}(e') \vee \text{tail}(e) = \text{tail}(e')].$$

Each such identification is called an *edge-folding* and we say that the edge e (as well as e') was *folded*. Figure 1 illustrates the process, which terminates in finitely many steps yielding the folding Γ_H . It is easy to verify that the folding so obtained is well-defined, and moreover, is independent of the choice of generating set for H . It is not hard to see that the rank of H is precisely $|E_H| - |V_H| + 1$.




Figure 1: Constructing a folding from a rose.

Now we consider subgroups of F_2 : If $H \leq_{\text{f.g.}} F_2$ then Γ_H has vertices of undirected degree ≤ 4 , where by “undirected degree” $d = d_H(v)$ of a vertex v we mean the sum of the number of outgoing and incoming edges at v . Put $d_i(\Gamma_H) = |\{v \in V_H \mid d_H(v) = i\}|$, for $i = 1, 2, 3, 4$. Vertices of degree 3 may be classified into 4 types, denoted $C_a, C_b, C_{a^{-1}}, C_{b^{-1}}$, based on the labels of the incident edges (see figure 2). For each $x \in \{a^{\pm}, b^{\pm}\}$, we define $C_x(\Gamma_H)$ to be the number of degree 3 vertices of type C_x in Γ_H . The rank of H can be computed by the formula

$$\text{rank}(H) = d_4(\Gamma_H) + \frac{d_3(\Gamma_H)}{2} - \frac{d_1(\Gamma_H)}{2} + 1$$

The graph-theoretic approach to Hanna Neumann’s conjecture is based on the following key observation [16, 8]. Consider the product automaton $\Gamma_H \times \Gamma_K$, whose vertex set is $V_H \times V_K$ and two vertices (u_1, v_1) and (u_2, v_2) are connected by an edge labelled x iff both $(u_1, v_1) \in E_H$ and $(u_2, v_2) \in E_K$ have label x . Consider the connected component Δ of $\Gamma_H \times \Gamma_K$ which contains $(1_H, 1_K)$. The




Figure 2: Local structure of vertex labels in a folding.

core of Δ is obtained by repeated deletion of all vertices of degree 1 different from $(1_H, 1_K)$. It is not hard to see that the core of Δ is precisely $\Gamma_{H \cap K}$.

Many proofs of the Hanna Neumann conjecture (for groups of particular ranks) require a case-by-case analysis based on the numbers and types of degree 3 vertices present in the foldings of H, K . The next theorem has implications on the number of cases which need to be considered in such arguments; it is proved in section 3.

Theorem 3. *Let $F = F(X)$ be a non-abelian free group. There is a monomorphism $\phi_0 : F \rightarrow F_2$ into the free group of rank 2 such that: for any groups $H, K \leq F$, the foldings of $H^{\phi_0}, K^{\phi_0} \leq_{f.g.} F_2$ have the property that all their branch vertices are of degree 3, and all degree 3 vertices have the same type.*

The previous theorem has the following immediate corollary:

Corollary 1. *If the Hanna Neumann conjecture is false then there is a counterexample $H, K \leq_{f.g.} F_2$ having the additional property that all the branch vertices in the foldings of H and K are of degree 3, and all degree 3 vertices have the same type.*

2 Preliminaries

The numbers $C_x(\Gamma_H)$ and $C_x(\Gamma_K)$ allow one to compute upper bounds on the numbers of vertices of degree 3 in $\Gamma_H \times \Gamma_K$ and hence in $\Gamma_{H \cap K}$. Observe that by considering a suitable conjugate of H and K we can always assume $d_1(\Gamma_H) = d_1(\Gamma_K) = 0$. Furthermore,

$$\begin{aligned} d_4(\Gamma_{H \cap K}) &\leq d_4(\Gamma_H \times \Gamma_K) = d_4(\Gamma_H)d_4(\Gamma_K), \text{ and} \\ d_3(\Gamma_{H \cap K}) &\leq d_3(\Gamma_H \times \Gamma_K) \\ &= d_4(\Gamma_H)d_3(\Gamma_K) + d_3(\Gamma_H)d_4(\Gamma_K) + \sum_{x \in \{a, b\}^\pm} C_x(\Gamma_H)C_x(\Gamma_K) \end{aligned}$$

Definition 1. Given two subgroups $H, K \leq_{f.g.} F_2$ and $x \in \{a, b\}^\pm$, we define

$$\begin{aligned} \delta_x(H, K) &= \min \left\{ \frac{C_x(\Gamma_H)}{d_3(\Gamma_H)}, \frac{C_x(\Gamma_K)}{d_3(\Gamma_K)} \right\} \\ \delta(H, K) &= \max_{x \in \{a^\pm, b^\pm\}} \delta_x(H, K) \end{aligned}$$

and put $\mu(H, K)$ to be any $x \in \{a^\pm, b^\pm\}$ for which $\delta_x(H, K) = \delta(H, K)$.

Remark 1. Walter Neumann [13] showed that if $H, K \leq_{f.g.} F_2$ are a counterexample to the conjecture, then $\delta(H, K) > \frac{1}{2}$. We outline his argument here, in graph-theoretic notation. Using a simple and beautiful argument from convexity theory, he showed that if $\delta(H, K) \leq \frac{1}{2}$ then

$$\sum_{x \in \{a, b\}^\pm} C_x(\Gamma_H) C_x(\Gamma_K) \leq \frac{1}{2} d_3(\Gamma_H) d_3(\Gamma_K).$$

It follows then that

$$\begin{aligned} \text{rank}(H \cap K) - 1 &= d_4(\Gamma_{H \cap K}) + d_3(\Gamma_{H \cap K})/2 - d_1(\Gamma_{H \cap K})/2 \\ &\leq d_4(\Gamma_{H \cap K}) + d_3(\Gamma_{H \cap K})/2 \\ &\leq d_4(\Gamma_H) d_4(\Gamma_K) + \frac{1}{2} \left[d_4(\Gamma_H) d_3(\Gamma_K) + d_3(\Gamma_H) d_4(\Gamma_K) \right. \\ &\quad \left. + \sum_{x \in \{a, b\}^\pm} C_x(\Gamma_H) C_x(\Gamma_K) \right] \\ &\leq d_4(\Gamma_H) d_4(\Gamma_K) + d_4(\Gamma_H) d_3(\Gamma_K)/2 + d_3(\Gamma_H) d_4(\Gamma_K)/2 \\ &\quad + d_3(\Gamma_H) d_3(\Gamma_K)/4 \\ &\leq [\text{rank}(H) - 1][\text{rank}(K) - 1]. \end{aligned}$$

and thus the conjecture holds.

3 Results

Remark 2. Given an endomorphism $\phi : F_2 \rightarrow F_2$, the folding Γ_{H^ϕ} can be obtained from Γ_H as follows. First construct a labelled directed graph $\phi(\Gamma_H)$ by replacing each edge with label x in Γ_H by a sequence of edges labelled by the successive letters of x^ϕ (for $x = a, b$). Then, apply the previously described folding procedure to transform the graph $\phi(\Gamma_H) \rightsquigarrow \Gamma_{H^\phi}$. One may verify that this yields a folding which is isomorphic to the one obtained by constructing Γ_{H^ϕ} directly from the set $\{w_1^\phi, \dots, w_n^\phi\}$.

For example, if ϕ is a length-preserving automorphism, then Γ_{H^ϕ} can be obtained from Γ_H by replacing every label x by x^ϕ and changing the orientation of the edges if necessary.

Lemma 1. *Let Γ_H be the folding of a subgroup $H \leq_{f.g.} F_2$, and $\phi : F_2 \rightarrow F_2$ an endomorphism. If two edges e, f from $\phi(\Gamma_H)$ get folded during the folding process $\phi(\Gamma_H) \rightsquigarrow \Gamma_{H^\phi}$, then there must exist a path p in $\phi(\Gamma_H)$ beginning at e and ending at f with the property that every edge in p was folded during the folding process.*

Proof. The lemma is proved by induction on the number n of edge-foldings which take place during the folding process—note that this number does not depend on the folding process since it is equal to $|E_{\phi(\Gamma_H)}| - |E_{\Gamma_{H^\phi}}|$ and the resultant folded graph Γ_{H^ϕ} is unique). For $n = 1$, the path p consists of just edges e, f . Now suppose the first edge-folding occurs when edges d_1 and d_2 are merged into an edge d' , and denote the folding obtained after this identification as Γ' . By induction, there exists a path p' in Γ' connecting e and f . There are two cases to consider: either d' appears in p' , or it does not. In the first case, let p_1 (resp. p_2) be the path obtained by replacing d' with d_1 (resp d_2) in p' . It is clear that either p_1 or p_2 must fulfill the requirements of the lemma. In the second case, we simply take $p = p'$. $\square$

Lemma 2. *Let $H, K \leq_{f.g.} F_2$ be subgroups which satisfy $\delta(H, K) > \frac{1}{2}$ (and hence are a potential counterexample to the Hanna Neumann conjecture) and take $*$, $\circ$ to be two length-preserving automorphisms of F_2 whose values differ on $\mu(H, K)$, i.e. $\mu(H, K)^* \neq \mu(H, K)^\circ$. Then the groups H°, K^* must satisfy the Hanna Neumann conjecture.*

Proof. Set $x_0 = \mu(H, K)$. Since $\delta(H, K) > \frac{1}{2}$, it follows that

$$\begin{aligned} C_{x_0}(\Gamma_H) &> \frac{1}{2}d_3(\Gamma_H) \\ C_{x_0}(\Gamma_K) &> \frac{1}{2}d_3(\Gamma_K). \end{aligned}$$

In light of Remark 2, Γ_{K^*} is the same graph as Γ_K , except that all a edges have been relabelled as a^* , and b edges have been relabelled as b^* , and an analogous statement is true about the relationship between Γ_{H° and Γ_H . So

$$\begin{aligned} C_x(\Gamma_K) &= C_{x^*}(\Gamma_{K^*}) \\ C_x(\Gamma_H) &= C_{x^\circ}(\Gamma_{H^\circ}) \end{aligned}$$

for all $x \in \{a^\pm, b^\pm\}$. It follows that

$$C_{x_0^*}(\Gamma_{K^*}) = C_{x_0}(\Gamma_K) > \frac{1}{2}d_3(\Gamma_K) = \frac{1}{2}d_3(\Gamma_{K^*}), \quad (2)$$

$$C_{x_0^\circ}(\Gamma_{H^\circ}) = C_{x_0}(\Gamma_H) > \frac{1}{2}d_3(\Gamma_H) = \frac{1}{2}d_3(\Gamma_{H^\circ}). \quad (3)$$

Since $x_0^\circ \neq x_0^*$, it follows that $\delta_x(H^\circ, K^*) < \frac{1}{2}$ for every $x \in \{a, b\}^\pm$. Thus, $\delta(H^\circ, K^*) < \frac{1}{2}$, and hence by Remark 1 the groups H°, K^* cannot be a counterexample to the conjecture. $\square$

The proofs of **Theorems 1 and 2** (see page 2) now follow from Lemma 2.

Proof. (Theorem 1) Suppose H, K do not satisfy the conjecture. By remark 1 we have $\delta(H, K) > \frac{1}{2}$. By definition of τ we have $\delta(H^\tau, K) = \delta(H, K)$. We

apply Lemma 2 to H^τ, K , taking $\circ$ to be the fixed-point-free length-preserving automorphism σ , and $*$ to be the identity automorphism. The theorem follows. $\square$

Proof. (Theorem 2) Suppose $X = \{a_1, \dots, a_n\}$. We consider the embedding $\psi : F(X) \rightarrow F_2 = F(a_1, a_2)$ defined by $\psi : a_i \mapsto a_1^i a_2 a_1^i$. If $H, K \leq_{f.g.} F(X)$ are a counterexample to the conjecture, then so are $H^\psi, K^\psi \leq_{f.g.} F_2$. Let $^-$ be the automorphism of $F(X)$ given by $a_i \mapsto a_i^{-1}$ (for each $a_i \in X$). Restricting $^-$ to F_2 and applying the previous lemma, we see that either H^ψ, K^ψ or $(H^\psi)^-, K^\psi$ must satisfy the conjecture. But $(H^-)^\psi = (H^\psi)^-$; here we think of H^ψ as a subgroup of $F(X)$ under the canonical inclusion of F_2 into $F(X)$. It follows that either H^ψ, K^ψ or $(H^-)^\psi, K^\psi$ must satisfy the conjecture. Since ψ is a monomorphism, this implies that either H, K or H^-, K must satisfy the conjecture. $\square$

Now towards the proof of Theorem 3, we introduce the following definition:

Definition 2. We say $\phi : F_2(a, b) \rightarrow F_2(a, b)$ is a N -endomorphism if it has the property that $U_\phi = \{a^\phi, b^\phi\}$ is N -reduced [10, pp.6], which is to say that every triple v_1, v_2, v_3 in $U_\phi^\pm$ satisfies

(N0) $v_1 \neq 1$,

(N1) $v_1 v_2 \neq 1$ implies $|v_1 v_2| \geq |v_1| + |v_2|$,

(N2) $v_1 v_2, v_2 v_3 \neq 1$ implies $|v_1 v_2 v_3| > |v_1| + |v_2| + |v_3|$.

Remark 3. It is well-known [10, pp.7] that if subset U of a free group F satisfies N0-N2, then one may associate with each $u \in U$ words $a(u), m(u) \in F$ with $m(u) \neq 1$ such that $u = a(u)m(u)a(u^{-1})^{-1}$ in F and having the property that for any $w = u_1 \cdots u_t$, $t \geq 0$, $u_i \in U^\pm$ where $u_i u_{i+1} \neq 1$, the subwords $m(u_1), \dots, m(u_t)$ remain uncanceled in the reduced form of w .

Lemma 3. Every N -endomorphism of F_2 is a monomorphism.

Proof. Take $w \in F_2$, with $w \neq 1$. By (N2), $|(w^\phi)^3| > 0$, hence $(w^\phi)^3 \neq 1$. It follows that $w^\phi \neq 1$. $\square$

Lemma 4. Given $H \leq_{f.g.} F_2$ and an N -endomorphism ϕ of F_2 , then for every edge e in Γ_H , at least one edge from the image of e under ϕ does not get folded during the folding process $\phi(\Gamma_H) \rightsquigarrow \Gamma_{H^\phi}$.

Proof. Let $e = (u, v)$ be any edge of Γ_H ; suppose e is labelled by $x \in \{a, b\}^\pm$. Consider the path $\phi(e)$ in $\phi(\Gamma_H)$; this path consists of a sequence of edges labelled by successive letters of x^ϕ . Since ϕ is an N -endomorphism, $\{a^\phi, b^\phi\}$ is N -reduced and Remark 3 applies. Accordingly, let $\bar{e}$ be the edge in $\phi(e)$ which corresponds to the first letter of $m(x^\phi)$ inside x^ϕ . We claim that $\bar{e}$ does not get folded during the folding process $\phi(\Gamma_H) \rightsquigarrow \Gamma_{H^\phi}$.

Suppose towards contradiction, that $\bar{e}$ gets folded with some edge f during the folding process $\phi(\Gamma_H) \rightsquigarrow \Gamma_{H^\phi}$. Then by Lemma 1, there must exist a non-backtracking path p in $\phi(\Gamma_H)$ beginning at $\bar{e}$ and ending at f , with the property that every edge in p was folded during the folding process. Since p is a non-backtracking path in $\phi(\Gamma_H)$, it is a subpath of $\phi(q)$ for some non-backtracking path q in Γ_H . It follows that the labels along $\phi(q)$ are a word $u_1 \cdots u_t$, $t \geq 0$, $u_i \in \{a^\phi, b^\phi\}^\pm$ and $u_i u_{i+1} \neq 1$. Since $\bar{e}$ is labelled by the first letter of $m(x^\phi)$, by Remark 3 the edge $\bar{e}$ was *not* folded during the folding process; this is a contradiction. $\square$

We introduce the following notations: let $\Gamma_H = (V_H, E_H)$ be the folding of H . Take any vertex $v \in V_H$, and let E_v be the edges incident to v . Define Γ_v to be the tree subgraph of Γ_H induced by edges E_v . Then $\phi(\Gamma_v)$ is also a tree.

By Lemma 4, we may associate to each edge $e \in E_v$, an edge $m(e) \in E_{\phi(\Gamma_H)}$ which does not get folded during the folding process $\phi(\Gamma_H) \rightsquigarrow \Gamma_{H^\phi}$. We define $\text{tr}\phi(\Gamma_v)$ to be the graph obtained by truncating the branches of $\phi(\Gamma_v)$ so that they terminate with edges $m(e)$, $e \in E_v$. Clearly, for all $v \in V_H$, $\text{tr}\phi(\Gamma_v)$ is a subgraph of $\phi(\Gamma_H)$.

The next lemma shows that N -endomorphisms do not cause large-scale disturbances in the neighborhood of branch vertices.

Lemma 5. *Given $H \leq_{f.g.} F_2$ and an N -endomorphism ϕ of F_2 , then during the folding process $\phi(\Gamma_H) \rightsquigarrow \Gamma_{H^\phi}$, no edge from $\text{tr}\phi(\Gamma_v)$ gets folded with an edge from outside $\text{tr}\phi(\Gamma_v)$.*

Proof. Suppose, towards contradiction, that an edge e inside $\text{tr}\phi(\Gamma_v)$ and an edge f outside $\text{tr}\phi(\Gamma_v)$ get folded during the folding process $\phi(\Gamma_H) \rightsquigarrow \Gamma_{H^\phi}$. Then by Lemma 1, there exists a path p beginning at e and ending at f with the property that every edge in p was folded during the folding process. Then p must pass through some edge $m(e)$, $e \in E_v$. This contradicts the properties of $m(e)$ as determined in Lemma 4. It follows that no edge inside $\text{tr}\phi(\Gamma_v)$ gets folded with an edge outside $\text{tr}\phi(\Gamma_v)$. $\square$

Informally stated, the previous lemma implies that for an N -endomorphism ϕ and subgroup $H \leq_{f.g.} F_2$, the 5-tuple of values

$$C_a(\Gamma_{H^\phi}), C_b(\Gamma_{H^\phi}), C_{a^{-1}}(\Gamma_{H^\phi}), C_{b^{-1}}(\Gamma_{H^\phi}), d_4(\Gamma_{H^\phi})$$

is completely determinable from the 5-tuple of values

$$C_a(\Gamma_H), C_b(\Gamma_H), C_{a^{-1}}(\Gamma_H), C_{b^{-1}}(\Gamma_H), d_4(\Gamma_H),$$

without knowledge of any further structure (e.g. the generating set) of H .

Lemma 6. *Let $\phi_0 : F_2 \rightarrow F_2$ be the endomorphism defined by $\phi_0 a = a^2$ and $\phi_0 b = [a, b]$. Then for any finitely generated subgroup $H \leq_{f.g.} F_2$, $C_{b^{-1}}(\Gamma_{H^{\phi_0}}) = d_3(\Gamma_{H^{\phi_0}})$, and $\text{rank}(H^{\phi_0}) = \text{rank}(H)$.*

Proof. It is straightforward to check that $\{a^2, [a, b]\}$ is N-reduced, and hence ϕ_0 is an N -endomorphism. By Lemma 5 it suffices to consider the effect of ϕ_0 on the various types of branch vertices in Γ_H . Figure 3 depicts how ϕ_0 transforms




Figure 3: The effect of ϕ_0 on a vertex of degree 4.

the neighborhood of degree 4 vertex v to produce two vertices of type C_{b-1} . The effect of ϕ_0 on each of the four types of degree 3 vertices may be determined simply by restricting our consideration to appropriate subgraphs of the depicted neighborhood. It follows that ϕ_0 transforms any degree 3 vertex in Γ_H into a vertex of type C_{b-1} in $\Gamma_{H^{\phi_0}}$. Thus,

$$C_{b-1}(\Gamma_{H^{\phi_0}}) = d_3(\Gamma_H) + 2d_4(\Gamma_H).$$

Since all branch vertices of Γ_H are seen to produce vertices of type C_{b-1} , we get that $C_{b-1}(\Gamma_{H^{\phi_0}}) = d_3(\Gamma_{H^{\phi_0}})$. Finally, by Lemma 3, ϕ_0 is a monomorphism, so $\text{rank}(H^{\phi_0}) = \text{rank}(H)$. $\square$

The proof of **Theorem 3** (see page 4) now follows from Lemma 6.

Proof. (Theorem 3) Fix the embedding $\psi : F(X) \rightarrow F_2 = F(a, b)$ defined by $\psi : a_i \mapsto a^i b a^i$. Put $H' = (H^\psi)^{\phi_0}$ and $K' = (K^\psi)^{\phi_0}$, where ϕ_0 is the endomorphism of $F(a, b)$ defined by $a^{\phi_0} = a^2$ and $b^{\phi_0} = [a, b]$. Then by Lemma 6,

$$\begin{aligned} C_{b-1}(\Gamma_{H'}) &= \frac{1}{2}d_3(\Gamma_{H'}) \\ C_{b-1}(\Gamma_{K'}) &= \frac{1}{2}d_3(\Gamma_{K'}), \end{aligned}$$

and hence $\delta(H', K') = 1$. $\square$

Corollary 1 follows immediately since $\psi\phi_0$ is a monomorphism, and hence $\text{rank}(H) = \text{rank}(H')$, $\text{rank}(K) = \text{rank}(K')$.

4 Acknowledgements

The first two authors are grateful to the Mathematics department at the City University of New York Graduate Center for funding this work as part of the

their ongoing doctoral research. The second author thanks both the Center for Computational Sciences at the Naval Research Laboratory in Washington DC, and the Advanced Engineering and Sciences division of ITT Industries for their support of these endeavors.

References

- [1] R. G. Burns. On the intersection of finitely generated subgroups of a free group. *Mathe. Z.*, 119:121–130, 1971.
- [2] Warren Dicks. Equivalence of the strengthened Hanna Neumann conjecture and the amalgamated graph conjecture. *Invent. Math.*, 117:373–389, 1994.
- [3] Edward Formanek and Warren Dicks. The rank three case of the Hanna Neumann conjecture. *J. Group Theory*, 4:113–151, 2001.
- [4] S. M. Gersten. Intersections of finitely generated subgroups of free groups and resolutions of graphs. *Invent. Math.*, 71:567–591, 1983.
- [5] A. G. Howson. On the intersection of finitely generated free groups. *J. London Math. Soc.*, 29:428–434, 1954.
- [6] W. Imrich. On finitely generated subgroups of free groups. *Arch. Math.*, 28:21–24, 1977.
- [7] W. Imrich. Subgroup theorems and graphs. *Combinatorial Mathematics V*, Lecture notes in mathematics, vol. 622:1–27, Melbourne, 1983.
- [8] Ilya Kapovich and Alexei Miasnikov. Stallings foldings and the subgroup structure of free groups. *J. Algebra*, 248 no 2:608–668, 2002.
- [9] Bilal Khan. Positively generated subgroups of free groups and the Hanna Neumann conjecture. In *Contemporary Mathematics, volume 296*. R. Gilman et al. American Mathematical Society, 2002.
- [10] Roger C. Lyndon and Paul E. Schupp. *Combinatorial Group Theory*. Springer-Verlag, New York, 1977.
- [11] John Meakin and Pascal Weil. Subgroups of free groups: a contribution to the Hanna Neumann conjecture. *Geom. Dedicata*, to appear.
- [12] H. Neumann. On intersections of finitely generated subgroups of free groups. *Publ. Math., Debrecen*, 5:186–189, 1956.
- [13] W. Neumann. On the intersection of finitely generated free groups. In *Canberra 1989 (Lecture notes in Math., vol. 1456, pp 161-170)*. Springer-Verlag, 1990.
- [14] P. Nickolas. Intersections of finitely generated free groups. *Bull. of the Australian Math. Soc.*, 34:339–348, 1985.

- [15] B. Servatius. A short proof of a theorem of Burns'. *Mathe. Z.*, 184:133–137, 1983.
- [16] J. R. Stallings. Topology of finite graphs. *Invent. Math.*, 71:551–565, 1983.
- [17] G. Tardos. On the intersection of subgroups of a free group. *Invent. Math.*, 108:29–36, 1992.
- [18] G. Tardos. Toward the Hanna Neumann Conjecture using Dicks' method. *Invent. Math.*, 123:95–104, 1996.